



OFFICE OF NAVAL RESEARCH

Contract N00014-85-K-0772

Work Unit No. 4326-808

TECHNICAL REPORT NO. 137

Metallacarborane Complexes That Incorporate the Lanthanides. Synthesis,
Molecular Structure, and Spectroscopic Characterization of Dicarbollide Complexes
of Samarium and Ytterbium

by

Mark J. Manning, Carolyn Knobler, Rajesh Khattar, and M. Frederick Hawthorne*



Prepared for Publication

in

Inorganic Chemistry

University of California at Los Angeles
Department of Chemistry and Biochemistry
Los Angeles, California 90024

May 1, 1991

A-1

Reproduction in whole or in part is permitted for
any purpose of the United States Government

This document has been approved for public release and sale;
its distribution is unlimited

91-07106



REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE			Distribution list enclosed		
4. PERFORMING ORGANIZATION REPORT NUMBER(S) 137			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION UCLA Dept. of Chem. & Biochem.		6b. OFFICE SYMBOL (If applicable)		7a. NAME OF MONITORING ORGANIZATION Office of Naval Research	
6c. ADDRESS (City, State, and ZIP Code) 6115 Young Hall 405 Hilgard Avenue Los Angeles, CA 90024				7b. ADDRESS (City, State, and ZIP Code) Department of the Navy Arlington, VA 22217	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (If applicable)		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER	
8c. ADDRESS (City, State, and ZIP Code)		10. SOURCE OF FUNDING NUMBERS			
		PROGRAM ELEMENT NO		PROJECT NO	TASK NO
					WORK UNIT ACCESSION NO
11. TITLE (Include Security Classification) Metallacarborane Complexes That Incorporate the Lanthanides. Synthesis, Molecular Structure, and Spectroscopic Characterization of Dicarbollide Complexes of Samarium and Ytterbium					
12. PERSONAL AUTHOR(S) Mark J. Manning, Carolyn B. Knobler, Rajesh Khattar, and M. Frederick Hawthorne					
13a. TYPE OF REPORT Technical Report		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Year, Month, Day) May 1, 1991	
15. PAGE COUNT 15					
16. SUPPLEMENTARY NOTATION To be published in <u>Inorganic Chemistry</u>					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	Metallacarborane, dicarbollide, samarium, ytterbium, thallium, sandwich compound		
19. ABSTRACT (Continue on reverse if necessary and identify by block number)					
<p>The interaction of $\text{Na}_2[\text{nido-7,8-C}_2\text{B}_9\text{H}_{11}]$ (1) with LnI_2 ($\text{Ln} = \text{Sm}, \text{Yb}$) in THF affords a complex with the composition $\text{Ln}(\text{C}_2\text{B}_9\text{H}_{11})(\text{THF})_4$ ($\text{Ln} = \text{Sm}$ (2), Yb (3)). Both these complexes have been fully characterized by spectroscopic techniques, and the molecular structure of the DMF derivative of 3 has been established by a single-crystal X-ray diffraction study. The DMF derivative, $\text{Yb}(\text{C}_2\text{B}_9\text{H}_{11})(\text{DMF})_4$ (3b), crystallizes in the orthorhombic space group $Pbca$ with $a = 10.208$ (1) Å, $b = 17.005$ (3) Å, $c = 31.627$ (5) Å, $V = 5479$ Å³, and $Z = 8$. Data were collected on a modified Picker FACS-1 diffractometer at 25 °C using Mo Kα radiation, to a maximum $2\theta = 50^\circ$, giving 4841 unique reflections, and the structure was solved by statistical methods. The final discrepancy index was $R = 0.082$, $R_w = 0.084$ for 1424 independent reflections with $I > 2\sigma(I)$. The dicarbollide ligand is η^3-bound to the Yb ion, and four DMF molecules are coordinated to Yb through the oxygen atoms. Complex 2 is fluxional in solution. The fluxionality of complex 2 in solution has been monitored by variable-temperature ¹¹B NMR spectroscopy, and a dynamic process involving the Sm^{2+} ion and the dicarbollide ligand has been proposed. The reaction of <i>closo</i>-1,1,1,1-(THF)₄-1,2,3-$\text{LnC}_2\text{B}_9\text{H}_{11}$ ($\text{Ln} = \text{Sm}$ (2), Yb (3)) with $[\text{PPN}]^+[\text{closo-3,1,2-TiC}_2\text{B}_9\text{H}_{11}]^-$ in THF produces $[\text{3,3-(THF)}_2\text{-} \text{closo-3,3'-Ln(3,1,2-LnC}_2\text{B}_9\text{H}_{11})_2]^-[\text{PPN}]^+$ ($\text{Ln} = \text{Sm}$ (4), Yb (5)), which has been characterized spectroscopically. The molecular structure of 4 has been established by an X-ray diffraction study. The complex 4 crystallizes in the triclinic space group $P\bar{1}$ with $a = 8.9374$ (3) Å, $b = 17.8703$ (6) Å, $c = 18.4989$ (7) Å, $\alpha = 107.5402$ (9)°, $\beta = 91.085$ (1)°, $\gamma = 90.705$ (1)°, $V = 2816$ Å³, and $Z = 2$. Data were collected at 25 °C on a diffractometer equipped with a small Huber circle, using Mo Kα radiation, to</p>					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL M. F. Hawthorne			22b. TELEPHONE (Include Area Code) (213)825-7378		22c. OFFICE SYMBOL

Metallacarborane Complexes That Incorporate the Lanthanides. Synthesis, Molecular Structure, and Spectroscopic Characterization of Dicarborollide Complexes of Samarium and Ytterbium

Mark J. Manning, Carolyn B. Knobler, Rajesh Khattar, and M. Frederick Hawthorne*

Received May 8, 1990

The interaction of $\text{Na}_2[\text{nido-7,8-C}_2\text{B}_9\text{H}_{11}]$ (1) with LnI_2 ($\text{Ln} = \text{Sm}, \text{Yb}$) in THF affords a complex with the composition $\text{Ln}(\text{C}_2\text{B}_9\text{H}_{11})(\text{THF})_4$ ($\text{Ln} = \text{Sm}$ (2), Yb (3)). Both these complexes have been fully characterized by spectroscopic techniques, and the molecular structure of the DMF derivative of 3 has been established by a single-crystal X-ray diffraction study. The DMF derivative, $\text{Yb}(\text{C}_2\text{B}_9\text{H}_{11})(\text{DMF})_4$ (3b), crystallizes in the orthorhombic space group $Pbca$ with $a = 10.208$ (1) Å, $b = 17.005$ (3) Å, $c = 31.627$ (5) Å, $V = 5479$ Å³, and $Z = 8$. Data were collected on a modified Picker FACS-1 diffractometer at 25 °C using Mo K α radiation, to a maximum $2\theta = 50^\circ$, giving 4841 unique reflections, and the structure was solved by statistical methods. The final discrepancy index was $R = 0.082$, $R_w = 0.084$ for 1424 independent reflections with $I > 2\sigma(I)$. The dicarborollide ligand is η^3 -bound to the Yb ion, and four DMF molecules are coordinated to Yb through the oxygen atoms. Complex 2 is fluxional in solution. The fluxionality of complex 2 in solution has been monitored by variable-temperature ¹¹B NMR spectroscopy, and a dynamic process involving the Sm^{2+} ion and the dicarborollide ligand has been proposed. The reaction of *closo*-1,1,1,1-(THF)₄-1,2,3- $\text{LnC}_2\text{B}_9\text{H}_{11}$ ($\text{Ln} = \text{Sm}$ (2), Yb (3)) with $[\text{PPN}]^+[\text{closo-3,1,2-TlC}_2\text{B}_9\text{H}_{11}]^-$ in THF produces $\{3,3-(\text{THF})_2\text{-}com\text{-}mo\text{-}3,3'\text{-Ln}(3,1,2\text{-}\text{LnC}_2\text{B}_9\text{H}_{11})_2\}^+[\text{PPN}]^-$ ($\text{Ln} = \text{Sm}$ (4), Yb (5)), which has been characterized spectroscopically. The molecular structure of 4 has been established by an X-ray diffraction study. The complex 4 crystallizes in the triclinic space group $P\bar{1}$ with $a = 8.9374$ (3) Å, $b = 17.8703$ (6) Å, $c = 18.4989$ (7) Å, $\alpha = 107.5402$ (9)°, $\beta = 91.085$ (1)°, $\gamma = 90.705$ (1)°, $V = 2816$ Å³, and $Z = 2$. Data were collected at 25 °C on a diffractometer equipped with a small Huber circle, using Mo K α radiation, to a maximum of $2\theta = 45^\circ$, giving 7358 unique reflections, and the structure was solved by a combination of conventional Patterson, Fourier, and full-matrix least-squares techniques. The final discrepancy index was $R = 0.053$, $R_w = 0.065$ for 5324 independent reflections with $I > 3\sigma(I)$. Both the dicarborollide ligands are η^3 -bound to the Sm ion in a nonparallel or bent fashion, and the coordination sphere about the Sm is completed by two THF molecules. The coordination geometry of 4 can be best described as a distorted tetrahedron. This is the first structurally authenticated example of a bis(dicarborollide)lanthanide complex. The structure of 4 is compared with related bis(cyclopentadienyl)lanthanides.

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1